

Novel construction and monodromy relation for 3pt-point functions @ weak coupling

Takuya Nishimura

Institute of physics, University of Tokyo, Komaba

June, 19th, 2015@Integrable Approaches to 3pt functions in
AdS₅/CFT₄

Based on

- Y. Kazama, S. Komatsu and T.N, [[arXiv:1410.8533](#)]
- Y. Kazama, S. Komatsu and T.N, [[arXiv:1506.03203](#)]

Introduction

Why 3pt functions in $\mathcal{N} = 4$ SYM?

- This is a workshop of the three-point functions.
- They are fundamental building blocks of the theory together with the 2pt. functions.
- They encode the dynamics of the string theory on the AdS background.

We need to study these fundamental observables in detail to reveal the underlying mechanism of AdS/CFT.

- Spectrum problem has been studied using various integrability techniques.
⇒ It means the underlying 1+1 dim system (light-cone gauge fixed sigma model, spin chain.) is integrable.
- However, “integrability” is not obvious beyond the spectrum problem a priori.

Is there any notion of “integrability” beyond the spectrum problem? If there is, what is the precise meaning of “integrability”?

Monodromy relation @strong coupling

At the strong coupling regime, the so-called monodromy relation plays an quite important role [’11 Janik,Wereszczynski],[’11,’12,’13 Kazama,Komatsu].

$$\Omega_1(u)\Omega_2(u)\Omega_3(u) = 1$$

- The monodromy relation provide a global information even without knowing the exact form of the vertex operators and the saddle pt. configuration.
- Combined with the analyticity, it determines the semi-classical three-point functions completely.

What is the weak coupling counter part of this relation? How is it useful to constrain three-point functions?

Motivations for our works

- 1 We would like to build weak coupling correlators **respecting the symmetry**.
- 2 We wish to find the weak coupling analogue of the monodromy relation.

Results

- 1 We develop a new formalism in which the symmetry is manifest.
⇒ We can simplify the analysis exploiting the symmetry.
- 2 We also derive the monodromy relations for correlator at weak coupling.


Plan of talk

- ① Introduction
- ② Construction of 3pt. functions @weak coupling
- ③ Monodromy relations
- ④ Summary and prospects

Construction of 3pt. functions @weak coupling

Tree-level three-point functions

Tree-level three-point functions are calculated by taking all possible **Wick contractions**. In particular, only **planar graphs** contribute in the large N_c limit.



- At tree-level, the dimensions of operators are highly **degenerate**.
- According to the usual **degenerate perturbation theory**, $\mathcal{O}_i(x_1), \mathcal{O}_j(x_2), \mathcal{O}_k(x_3)$ must be eigenstates of the 1-loop dilatation operator.

⇒ **Combinatorics of Wick contractions.**

The tailoring gives an efficient method to calculate relevant contractions
[’11 Escobedo, Gromov, Sever, Vieira].

- 1 Composite operators are mapped to eigenstates of the spin chain Hamiltonian $|\Psi_i\rangle$ [’02 Minahan, Zarembo, ’02 Beisert].
- 2 **Cutting** spin chains into the subchains: $|\Psi_i\rangle = \sum |\Psi_i\rangle_l \otimes |\Psi_i\rangle_r$.
- 3 **Flipping** the half of the states: $|\Psi_i\rangle \rightarrow \hat{\Psi}_i = \sum |\Psi_i\rangle_l \otimes_r \overleftarrow{\langle \Psi_i |}$
- 4 **Sewing** the states: $C_{ijk} \propto \sum \sum \sum \overleftarrow{\langle \Psi_i | \Psi_j \rangle} \overleftarrow{\langle \Psi_j | \Psi_k \rangle} \overleftarrow{\langle \Psi_k | \Psi_i \rangle}$

Let us consider the SU(2) sector, where the operators are composed of two types of scalar fields, say, $Z = \phi_1 + i\phi_2$, $X = \phi_3 + i\phi_4$.

- $Z \leftrightarrow |\uparrow\rangle$, $X \leftrightarrow |\downarrow\rangle$.
- Each operator is identified with an eigenstate of the XXX Heisenberg spin chain (**Bethe state**).

$$\mathcal{O}(x) \leftrightarrow |\mathbf{u}\rangle = \prod_{i=1}^M B(u_i) |\uparrow^\ell\rangle, \quad \Omega(u) = \begin{pmatrix} A(u) & B(u) \\ C(u) & D(u) \end{pmatrix}$$

$$\Omega(u) = L_1(u) \cdots L_\ell(u), \quad L_n(u) = \begin{pmatrix} u + iS_n^3 & iS_n^- \\ iS_n^+ & u - iS_n^3 \end{pmatrix}$$

$$\prod_{k=1}^{\ell} \left(\frac{u_k + i/2}{u_k - i/2} \right)^{\ell} = \prod_{j \neq k} \frac{u_k - u_j + i}{u_k - u_j - i}$$

A special class of 3pt. functions in SU(2) sector

$$\mathcal{O}_1 = \sum \text{Tr}[ZX \cdots] \leftrightarrow \sum |\uparrow\downarrow \cdots\rangle$$

$$\mathcal{O}_2 = \sum \text{Tr}[\bar{Z}\bar{X} \cdots] \leftrightarrow \sum |\uparrow\downarrow \cdots\rangle$$

$$\mathcal{O}_3 = \sum \text{Tr}[Z\bar{X} \cdots] \leftrightarrow \sum |\uparrow\downarrow \cdots\rangle$$

The tailoring gives the structure constant in terms of scalar product of Bethe states.

Remarks

- In this special case, the result only involves $\langle \text{off-shell} | \text{on-shell} \rangle$.
 \Rightarrow Determinant formulas.
- The mapping to a single SU(2) spin chain is not natural from a symmetry point of view.

$SU(2)_L \times SU(2)_R$ double spin chain formalism

From a symmetry point of view, it is natural to introduce a **double spin chain** ['15 Kazama, Komatsu, T.N].

$$\Phi_{a\tilde{a}} = \begin{pmatrix} Z & X \\ -\bar{X} & \bar{Z} \end{pmatrix}_{a\tilde{a}} \leftrightarrow \begin{pmatrix} |\uparrow\rangle_L \otimes |\uparrow\rangle_R & |\uparrow\rangle_L \otimes |\downarrow\rangle_R \\ |\downarrow\rangle_L \otimes |\uparrow\rangle_R & |\downarrow\rangle_L \otimes |\downarrow\rangle_R \end{pmatrix}_{a\tilde{a}}$$

- $SO(4) \cong SU(2)_L \times SU(2)_R$ transformation is $\Phi \rightarrow g_L \Phi g_R$.
- A general $SO(4)$ scalar is labeled by bi-spinor $P^{a\tilde{a}}$: $P \cdot \Phi = P^{a\tilde{a}} \Phi_{a\tilde{a}}$.

$$P \cdot \Phi \leftrightarrow P^{1\tilde{1}} |\uparrow\rangle_L \otimes |\uparrow\rangle_R + P^{1\tilde{2}} |\uparrow\rangle_L \otimes |\downarrow\rangle_R \\ + P^{2\tilde{1}} |\downarrow\rangle_L \otimes |\uparrow\rangle_R + P^{2\tilde{2}} |\downarrow\rangle_L \otimes |\downarrow\rangle_R$$

General rotated vacua

To treat generic operators in the $SU(2)$ sector, we first consider rotated vacua on which non-BPS operators are build.

- General BPS operators are obtained by $SO(4)$ rotations

$$\text{Tr}[Z^\ell] \rightarrow \text{Tr}[(P \cdot \Phi)^\ell]:$$

$$Z = \Phi_{1\bar{1}} \rightarrow (g_L \Phi g_R)_{1\bar{1}} = (g_L)_1^a \Phi_{a\bar{a}} (g_R)^{\bar{a}}_{\bar{1}},$$
$$P^{a\bar{a}} = \mathbf{n}^a \tilde{\mathbf{n}}^{\bar{a}}, \mathbf{n}^a = (g_L)_1^a, \tilde{\mathbf{n}}^{\bar{a}} = (g_R)^{\bar{a}}_{\bar{1}}$$

- Hence, the vacua are assigned two **polarization spinors**:

$$\text{Tr}[(P \cdot \Phi)^\ell] \leftrightarrow |\mathbf{n}^\ell\rangle_L \otimes |\tilde{\mathbf{n}}^\ell\rangle_R,$$

$$|\mathbf{n}^\ell\rangle_L = |\mathbf{n}\rangle_L \otimes \cdots \otimes |\mathbf{n}\rangle_L, |\tilde{\mathbf{n}}^\ell\rangle_R = |\tilde{\mathbf{n}}\rangle_R \otimes \cdots \otimes |\tilde{\mathbf{n}}\rangle_R,$$

$$|\mathbf{n}\rangle_L = \mathbf{n}^1 |\uparrow\rangle_L + \mathbf{n}^2 |\downarrow\rangle_L, |\tilde{\mathbf{n}}\rangle_R = \tilde{\mathbf{n}}^{\bar{1}} |\uparrow\rangle_R + \tilde{\mathbf{n}}^{\bar{2}} |\downarrow\rangle_R.$$

- For convenience, we normalize polarization spinors as

$$\mathbf{n}^a \bar{\mathbf{n}}_a = \tilde{\mathbf{n}}^{\bar{a}} \tilde{\bar{\mathbf{n}}}_{\bar{a}} = 1, \bar{\mathbf{n}}_a = (\mathbf{n}^a)^*, \tilde{\bar{\mathbf{n}}}_{\bar{a}} = (\tilde{\mathbf{n}}^{\bar{a}})^*$$

General non-BPS operators in SU(2) sector

Non-BPS operators in the SU(2) sector can be obtained by adding magnons either on the SU(2)_L or SU(2)_R sector:

Type I: $|\mathbf{u}; \mathbf{n}^\ell\rangle_L \otimes |\tilde{\mathbf{n}}^\ell\rangle_R$, Type II: $|\mathbf{n}^\ell\rangle_L \otimes |\tilde{\mathbf{u}}; \tilde{\mathbf{n}}^\ell\rangle_R$.

Such states are obtained by SU(2)_L × SU(2)_R transformation. For example,

$$|\mathbf{u}; \mathbf{n}^\ell\rangle_L = g_L |\mathbf{u}; \uparrow^\ell\rangle_L, \quad |\mathbf{n}^\ell\rangle = g_L |\uparrow^\ell\rangle, \quad g_L \in \text{SU}(2)/\text{U}(1),$$

where $|\mathbf{u}; \uparrow^\ell\rangle_L = \prod_{i=1}^M B_L(u_i) |\uparrow^\ell\rangle_L$. It is convenient to parametrize g_L as

$$g_L = e^{\zeta S_- - \bar{\zeta} S_+} = e^{z S_-} e^{-\ln(1+|z|^2) S_3} e^{-\bar{z} S_+}$$

where $z = \frac{\zeta}{|\zeta|} \tan |\zeta|$. This is the so-called **coherent state representation** and z is the **projective coordinate** for the coset $\text{SU}(2)/\text{U}(1) \equiv S^2$.

With this parametrization $g_L = e^{zS_-} e^{-\ln(1+|z|^2)S_3} e^{-\bar{z}S_+}$, we find

$$|\mathbf{n}\rangle_L = g_L |\uparrow\rangle_L, \quad \mathbf{n} = g_L \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \frac{1}{\sqrt{1+|z|^2}} \begin{pmatrix} 1 \\ z \end{pmatrix},$$

Furthermore, the Bethe states on the rotated vacuum are expressed as follows

$$|\mathbf{u}; \mathbf{n}^\ell\rangle_L = g_L |\mathbf{u}; \uparrow^\ell\rangle_L = \left(\frac{1}{1+|z|^2} \right)^{\ell/2-M} e^{zS_-} |\mathbf{u}; \uparrow^\ell\rangle_L.$$

Notice that the Bethe state is a **highest weight state** of $SU(2)_L$, i.e. $S_+ |\mathbf{u}; \uparrow^\ell\rangle_L = 0$. Similarly, we have

$$|\mathbf{u}; \tilde{\mathbf{n}}^\ell\rangle_R = g_R |\tilde{\mathbf{u}}; \uparrow^\ell\rangle_R = \left(\frac{1}{1+|\tilde{z}|^2} \right)^{\ell/2-\tilde{M}} e^{\tilde{z}\tilde{S}_-} |\tilde{\mathbf{u}}; \uparrow^\ell\rangle_R.$$

Wick contraction as singlet projection

The Wick contractions for Z, X and their conjugate are summarized as

$$\overline{\Phi_{a\tilde{a}}\Phi_{b\tilde{b}}} = \epsilon_{ab}\epsilon_{\tilde{a}\tilde{b}}, \quad \Phi_{a\tilde{a}} = \begin{pmatrix} Z & X \\ -\bar{X} & \bar{Z} \end{pmatrix}_{a\tilde{a}}$$

To implement the above contractions in a spin chain language, we introduce a **singlet projector**

$$\begin{aligned} \langle \mathbf{1} | &= \epsilon_{ab} \langle a | \otimes \langle b |, \\ | \mathbf{1} \rangle &= | \uparrow \rangle, \quad | \mathbf{2} \rangle = | \downarrow \rangle, \quad \langle a | b \rangle = \delta_{ab}. \end{aligned}$$

With the singlet projector, we find

$$\overline{\Phi_{a\tilde{a}}\Phi_{b\tilde{b}}} = {}_L \langle \mathbf{1} | (|a\rangle_L \otimes |b\rangle_L) \cdot {}_R \langle \mathbf{1} | (|a\rangle_R \otimes |b\rangle_R)$$

For $P_1 \cdot \Phi = \mathbf{n}_1^a \tilde{\mathbf{n}}_1^{\tilde{a}} \Phi_{a\tilde{a}}$ and $P_2 \cdot \Phi = \mathbf{n}_2^b \tilde{\mathbf{n}}_2^{\tilde{b}} \Phi_{b\tilde{b}}$,

$$\begin{aligned} P_1 \cdot \overline{\Phi P_2 \cdot \Phi} &= {}_L \langle \mathbf{1} | (|\mathbf{n}_1\rangle_L \otimes |\mathbf{n}_2\rangle_L) \cdot {}_R \langle \mathbf{1} | (|\tilde{\mathbf{n}}_1\rangle_R \otimes |\tilde{\mathbf{n}}_2\rangle_R) \\ &= \langle \mathbf{n}_1, \mathbf{n}_2 \rangle \langle \tilde{\mathbf{n}}_1, \tilde{\mathbf{n}}_2 \rangle = (\epsilon_{ab} \mathbf{n}_1^a \mathbf{n}_2^b) (\epsilon_{\tilde{a}\tilde{b}} \tilde{\mathbf{n}}_1^{\tilde{a}} \tilde{\mathbf{n}}_2^{\tilde{b}}) \end{aligned}$$


Composite operators in the $SU(2)$ sector are schematically written as

$$\mathcal{O}_i \mapsto |\mathcal{O}_i\rangle_L \otimes |\tilde{\mathcal{O}}_i\rangle_R .$$

The Wick contractions for the composite operators are given by

$$\overline{\mathcal{O}_1 \mathcal{O}_2} = \langle |\mathcal{O}_1\rangle_L, |\mathcal{O}_2\rangle_L \rangle \cdot \langle |\tilde{\mathcal{O}}_1\rangle_R, |\tilde{\mathcal{O}}_2\rangle_R \rangle$$

$$\langle |\Psi_1\rangle, |\Psi_2\rangle \rangle = \langle \mathbf{1}_{12} | (|\Psi_1\rangle \otimes |\Psi_2\rangle) \rangle, \quad \langle \mathbf{1}_{12} | = \prod_{k=1}^{\ell} \langle \mathbf{1}_{k, \ell-k+1} |$$



The contributions from the left and right sector are completely *factorized*.

Cutting and sewing in our formalism

Let us concentrate on the $SU(2)_L$ part.


① Cutting:

$$|\mathcal{O}_i\rangle_L \rightarrow \sum_a |\mathcal{O}_{i_a}\rangle^l \otimes |\mathcal{O}_{i_a}\rangle^r \quad (i = 1, 2, 3)$$

② Sewing:

$$\begin{aligned} & \langle |\mathcal{O}_1\rangle_L, |\mathcal{O}_2\rangle_L, |\mathcal{O}_3\rangle_L \rangle \\ &= \sum_{a,b,c} \langle |\mathcal{O}_{1_a}\rangle^r, |\mathcal{O}_{2_b}\rangle^l \rangle \langle |\mathcal{O}_{2_b}\rangle^r, |\mathcal{O}_{3_c}\rangle^l \rangle \langle |\mathcal{O}_{3_c}\rangle^r, |\mathcal{O}_{1_a}\rangle^l \rangle \end{aligned}$$

$$C_{123} = \frac{\sqrt{l_1 l_2 l_3}}{N_c \sqrt{\mathcal{N}_1 \mathcal{N}_2 \mathcal{N}_3}} \langle |\mathcal{O}_1\rangle_L, |\mathcal{O}_2\rangle_L, |\mathcal{O}_3\rangle_L \rangle \cdot \langle |\tilde{\mathcal{O}}_1\rangle_R, |\tilde{\mathcal{O}}_2\rangle_R, |\tilde{\mathcal{O}}_3\rangle_R \rangle$$



z dependence from the Ward identities

Three-point functions must satisfy the Ward identity for $SU(2)_L$:

$$0 = \langle S^* | \mathcal{O}_1 \rangle_L, | \mathcal{O}_2 \rangle_L, | \mathcal{O}_3 \rangle_L \rangle + \langle | \mathcal{O}_1 \rangle_L, S^* | \mathcal{O}_2 \rangle_L, | \mathcal{O}_3 \rangle_L \rangle \\ + \langle | \mathcal{O}_1 \rangle_L, | \mathcal{O}_2 \rangle_L, S^* | \mathcal{O}_3 \rangle_L \rangle, \quad S^* \in \mathfrak{su}(2)_L.$$

Putting $|\hat{\mathcal{O}}_i\rangle_L = e^{z_i S^-} |\mathbf{u}_i; \uparrow^\ell\rangle$ into the above, we obtain the following **differential representation** for the Ward identities:

$$\sum_{i=1}^3 \rho_{z_i}(S^*) \langle | \hat{\mathcal{O}}_1 \rangle_L, | \hat{\mathcal{O}}_2 \rangle_L, | \hat{\mathcal{O}}_3 \rangle_L \rangle = 0,$$

$$\rho_{z_i}(S^-) = \frac{d}{dz_i}, \quad \rho_{z_i}(S^3) = L_i - z_i \frac{d}{dz_i}, \quad \rho_{z_i}(S^+) = L_i z_i - \frac{z_i^2}{2} \frac{d}{dz_i}.$$

L_i : $\mathfrak{su}(2)_L$ charge. These equations completely fix the z_i dependence:

$$\langle | \hat{\mathcal{O}}_1 \rangle_L, | \hat{\mathcal{O}}_2 \rangle_L, | \hat{\mathcal{O}}_3 \rangle_L \rangle = z_{21}^{L_1+L_2-L_3} z_{32}^{L_2+L_3-L_1} z_{13}^{L_3+L_1-L_2} \mathcal{G}$$

Representation in terms of pDWPF

The cutting for the Bethe state $|\mathbf{u}; \uparrow^\ell\rangle$ is determined in EGSV:

$$|\mathbf{u}; \uparrow^\ell\rangle \rightarrow \sum_{\alpha_l \cup \alpha_r = \mathbf{u}} H_\ell(\alpha_l, \alpha_r) |\alpha_l; \uparrow^{\ell_l}\rangle \otimes |\alpha_r; \uparrow^{\ell_r}\rangle ,$$

$$H_\ell(\alpha_l, \alpha_r) = \prod_{u \in \alpha_l} \prod_{v \in \alpha_r} \left(u - \frac{i}{2}\right)^{\ell_r} \left(v + \frac{i}{2}\right)^{\ell_l} \left(\frac{u - v + i}{u - v}\right)$$

From $|\mathbf{u}_i; \mathbf{n}_i^{\ell_i}\rangle \propto e^{z_i S_-} |\mathbf{u}_i; \uparrow^{\ell_i}\rangle$, we find

$$|\mathbf{u}_i; \mathbf{n}_i^{\ell_i}\rangle \rightarrow \sum_{\alpha_l \cup \alpha_r = \mathbf{u}} H_\ell(\alpha_l, \alpha_r) e^{z_i S_-} |\alpha_l; \uparrow^{\ell_l}\rangle \otimes e^{z_i S_-} |\alpha_r; \uparrow^{\ell_r}\rangle .$$

The sewing procedure produces the following building blocks:

$$\left\langle e^{z_i S_-} |\mathbf{x}; \uparrow^\ell\rangle, e^{z_j S_-} |\mathbf{y}; \uparrow^\ell\rangle \right\rangle$$

- ① Using the defining property of the singlet $\langle \mathbf{1}_{12} | (S^{(1)} + S^{(2)}) = 0$,

$$\left\langle e^{z_i S_-} | \mathbf{x}; \uparrow^\ell \right\rangle, e^{z_j S_-} | \mathbf{y}; \uparrow^\ell \right\rangle = \left\langle | \mathbf{x}; \uparrow^\ell \right\rangle, e^{(z_j - z_i) S_-} | \mathbf{y}; \uparrow^\ell \right\rangle .$$

- ② With the relation $\langle \mathbf{1}_{12} | B^{(1)}(u) = -\langle \mathbf{1}_{12} | B^{(2)}(u)$, we can gather all the excitations to the one side:

$$(-1)^{M_i} \left\langle | \uparrow^\ell \right\rangle, e^{(z_j - z_i) S_-} | \mathbf{x} \cup \mathbf{y}; \uparrow^\ell \right\rangle$$

- ③ From $\langle \mathbf{1}_{12} | (| \uparrow^\ell \rangle \otimes | \Psi_2 \rangle) = \langle \downarrow^\ell | \Psi_2 \rangle$, one finds

$$(-1)^{M_i} \langle \downarrow^\ell | e^{(z_j - z_i) S_-} | \mathbf{x} \cup \mathbf{y}; \uparrow^\ell \rangle = (-1)^{M_i} (z_j - z_i)^{\ell - M_i - M_j} Z_\ell(\mathbf{x} \cup \mathbf{y})$$

where $Z_\ell(\mathbf{u})$ is the so-called **partial Domain Wall Partition Function (pDWPF)**:

$$Z_\ell(\mathbf{u}) = \frac{1}{(\ell - M)!} \langle \downarrow^\ell | (S_-)^{\ell - M} \prod_{i=1}^M B(u_i) | \uparrow^\ell \rangle .$$

Finally, we have

$$\begin{aligned}
 & \langle |\mathcal{O}_1\rangle_L, |\mathcal{O}_2\rangle_L, |\mathcal{O}_3\rangle_L \rangle \\
 &= \left(\frac{1}{1 + |z_1|^2} \right)^{\ell_1/2 - M_1} \left(\frac{1}{1 + |z_2|^2} \right)^{\ell_2/2 - M_2} \left(\frac{1}{1 + |z_3|^2} \right)^{\ell_3/2 - M_3} \\
 &\times \sum_{\alpha_l^{(k)} \cup \alpha_r^{(k)} = \mathbf{u}_k} z_{21}^{\ell_{12} - |\alpha_r^{(1)}| - |\alpha_l^{(2)}|} z_{32}^{\ell_{23} - |\alpha_r^{(2)}| - |\alpha_l^{(3)}|} z_{13}^{\ell_{31} - |\alpha_r^{(3)}| - |\alpha_l^{(1)}|} \mathcal{D}_{\{\alpha_{l,r}^{(1)}, \alpha_{l,r}^{(2)}, \alpha_{l,r}^{(3)}\}} \\
 &\mathcal{D}_{\{\alpha_{l,r}^{(1)}, \alpha_{l,r}^{(2)}, \alpha_{l,r}^{(3)}\}} \\
 &= (-1)^{|\alpha_r^{(1)}| + |\alpha_r^{(2)}| + |\alpha_r^{(3)}|} \prod_{k=1}^3 H_{\ell_k}(\alpha_l^{(k)}, \alpha_r^{(k)}) Z_{\ell_{kk+1}}(\alpha_r^{(k)} \cup \alpha_l^{(k+1)})
 \end{aligned}$$

where $z_{ij} := z_i - z_j$, $|\alpha|$ is the number of elements in α .

- pDWPF has a determinant expression.
- The summation should be simplified so that it reproduces the correct kinematical dependence.

Kinematical dependence and further simplification

In particular, if we consider $|\mathcal{O}_1\rangle_L = |\mathbf{u}; \mathbf{n}_1\rangle$, $|\mathcal{O}_2\rangle_L = |\mathbf{v}; \mathbf{n}_2\rangle$,
 $|\mathcal{O}_3\rangle_L = |\mathbf{n}_3\rangle$,

$$\langle |\hat{\mathcal{O}}_1\rangle_L, |\hat{\mathcal{O}}_2\rangle_L, |\hat{\mathcal{O}}_3\rangle_L \rangle = z_{12}^{\ell_{21}-M_1-M_2} z_{32}^{\ell_{23}-M_2+M_1} z_{13}^{\ell_{31}-M_1+M_2} \mathcal{G} ,$$

which yields the following highest order term in z_3 :

$$(-1)^{\ell_{31}-M_1+M_2} z_3^{\ell_3} z_{21}^{\ell_{12}-M_1-M_2} \mathcal{G} .$$

On the other hands, the higher order term in the summation we have derived is given by

$$(-1)^{\ell_{31}} z_3^{\ell_3} z_{21}^{\ell_{12}-M_1-M_2} \mathcal{D}_{\{\alpha_{l,r}^{(1)}, \alpha_{l,r}^{(2)}, \emptyset\}} \Big|_{\alpha_l^{(1)} = \alpha_r^{(2)} = \emptyset}$$

By comparing the z_3 dependence,

$$\mathcal{G} = (-1)^{M_1+M_2} \mathcal{D}_{\{\alpha_{l,r}^{(1)}, \alpha_{l,r}^{(2)}, \emptyset\}} \Big|_{\alpha_l^{(1)} = \alpha_r^{(2)} = \emptyset} \propto H(\emptyset, \mathbf{u}) H(\mathbf{v}, \emptyset) Z_{\ell_{12}}(\mathbf{u} \cup \mathbf{v})$$

The result is

$$\begin{aligned} \langle |\mathcal{O}_1\rangle_L, |\mathcal{O}_2\rangle_L, |\mathcal{O}_3\rangle_L \rangle &= z_{12}^{\ell_{21}-M_1-M_2} z_{32}^{\ell_{23}-M_2+M_1} z_{13}^{\ell_{31}-M_1+M_2} \\ &\times \left(\frac{1}{1+|z_1|^2} \right)^{\ell_1/2-M_1} \left(\frac{1}{1+|z_2|^2} \right)^{\ell_2/2-M_2} \left(\frac{1}{1+|z_3|^2} \right)^{\ell_3/2-M_3} \\ &\times \prod_{k=1}^{M_1} \left(u_k + \frac{i}{2} \right)^{\ell_{31}} \prod_{l=1}^{M_2} \left(v_l - \frac{i}{2} \right)^{\ell_{23}} Z_{\ell_{12}}(\mathbf{u} \cup \mathbf{v}) \end{aligned}$$

Comments

- pDWPF has a determinant expression and the semi-classical limit is obtained ['12, Kostov].
- We have treated correlators for two type I operators and one type II operator: $|\mathbf{u}_i; \mathbf{n}_i\rangle_L \otimes |\tilde{\mathbf{n}}_i\rangle_R$ ($i = 1, 2$), $|\mathbf{n}_3\rangle_L \otimes |\tilde{\mathbf{u}}_3; \tilde{\mathbf{n}}_3\rangle_R$.
 \Rightarrow **A special class of 3pt. functions in EGSV is contained in this class.**
- The correlators for three type I operators, namely, $|\mathbf{u}_i; \mathbf{n}_i\rangle_L \otimes |\tilde{\mathbf{n}}_i\rangle_R$ ($i = 1, 2, 3$) involves a sum over partitions, even after the simplification.

- ① The Wick contractions are efficiently performed using the **singlet projector** $\langle \mathbf{1}_{12} |$.
- ② The defining property of the singlet ensures the **Ward identities** of the symmetry.
- ③ In this formalism, it is possible to deal a more general class of 3pt. functions and the structure constants can be expressed in terms of **pDWPFs**, rather than the scalar products.
- ④ The Ward identities and the kinematical dependence greatly simplify the result.

Monodromy relations for correlators @weak coupling

Monodromy relation for basic 2pt. functions

It is sufficient to consider the 2pt. point functions. We would like to show the following form of monodromy relation

$$\langle (\Omega(u - i/2))_{ik} | \mathcal{O}_1 \rangle, (\Omega(u + i/2))_{kj} | \mathcal{O}_2 \rangle \rangle = f(u) \delta_{ij} \langle | \mathcal{O}_1 \rangle, | \mathcal{O}_2 \rangle \rangle ,$$
$$\Omega(u) = L_1(u) \cdots L_\ell(u) , \quad L_n(u) = \begin{pmatrix} u + iS_n^3 & iS_n^- \\ iS_n^+ & u - iS_n^3 \end{pmatrix}$$

To prove this, we use two important equations:

Crossing : $\langle \mathbf{1}_{12} | L_n^{(1)}(u) = -\langle \mathbf{1} | L_{\ell-n+1}^{(2)}(-u)$

Inversion : $L_n(-u + i/2)L_n(u + i/2) = -(u^2 + 1)\mathbf{1}$

Crossing relation

To prove the crossing relation, notice that $L_n^{(k)}(u) = u + i\vec{S}_n^{(k)} \cdot \vec{\sigma}$, where $\vec{\sigma}$'s are Pauli matrices.

$$\begin{aligned}\langle \mathbf{1}_{12} | L_n^{(1)}(u) &= \langle \mathbf{1}_{12} | (u + i\vec{S}_n^{(1)} \cdot \vec{\sigma}) = \langle \mathbf{1}_{12} | (u - i\vec{S}_{\ell-n+1}^{(2)} \cdot \vec{\sigma}) \\ &= -\langle \mathbf{1}_{12} | L_{\ell-n+1}^{(2)}(-u) = \sigma^2 L_{\ell-n+1}^{(2)T}(u) \sigma^2\end{aligned}$$

With the crossing relation for the Lax operators, we have

$$\begin{aligned}\langle \mathbf{1}_{12} | \Omega^{(1)}(u) &= (-1)^\ell \langle \mathbf{1}_{12} | \overleftarrow{\Omega}^{(2)}(-u) = \langle \mathbf{1}_{12} | \sigma^2 \Omega^{(2)T}(u) \sigma^2, \\ \overleftarrow{\Omega}(u) &:= L_\ell(u) \cdots L_1(u), \quad \sigma^2 \Omega^T(u) \sigma^2 = \begin{pmatrix} D(u) & -B(u) \\ -C(u) & A(u) \end{pmatrix}\end{aligned}$$

In particular, one finds $\langle \mathbf{1}_{12} | B^{(1)}(u) = -\langle \mathbf{1}_{12} | B^{(2)}(u)$.

Inversion relation

By the explicit calculation, we can show

$$L(v)L(u) = (vu - 3/4) + i(v + u - i)\vec{S} \cdot \vec{\sigma}$$

Hence,

$$\begin{aligned} & \langle \mathbf{1}_{12} | \Omega^{(1)}(u - i/2) \Omega^{(2)}(u + i/2) \\ &= (-1)^\ell \langle \mathbf{1}_{12} | \overleftarrow{\Omega}^{(2)}(-u + i/2) \Omega^{(2)}(u + i/2) \\ &= (-1)^\ell \langle \mathbf{1}_{12} | \cdots L_1(-u + i/2) L_1(u + i/2) \cdots \\ &= (u^2 + 1)^\ell \langle \mathbf{1}_{12} | \end{aligned}$$

- To invert the Lax operator, the **shift for the spectral parameter** is necessary.
- For the case of $SL(2)$, we obtain the similar result, with shift for the spectral parameter.
- These results are obtained by the reduction from monodromy relation of $\mathfrak{psu}(2, 2|4)$ sector [’14 Jiang, Kostov, Petrovskii, Serban], [’15 Kazama, Komatsu, T.N].

If we consider the limit $u \rightarrow \infty$ and expand the monodromy relation in power of $1/u$, we obtain the Ward identities of the form

$$\langle S^*|\mathcal{O}_1\rangle, |\mathcal{O}_2\rangle\rangle + \langle |\mathcal{O}_1\rangle, S^*|\mathcal{O}_2\rangle\rangle = 0 .$$

- The expansion of the monodromy matrix in power of $1/u$ generates the Yangian generators.
 \Rightarrow *The monodromy relations include a kind of the Ward identities of them.*

Monodromy relation for harmonic R-matrix

We can also derive the monodromy relation for the R-matrix whose auxiliary space is the singleton representation ['15 Kazama, Komatsu, T.N].

$$\mathbf{R}_{12}(u) = (-1)^{\mathbb{J}} \frac{\Gamma(\mathbb{J} + u + 1)}{\Gamma(\mathbb{J} - u + 1)} \in \text{End}(\mathcal{V} \otimes \mathcal{V}) , \quad \mathcal{C}_2 = \mathbb{J}(\mathbb{J} + 1)$$

$$\langle \mathbf{1}_{12} | \Omega^{(1)}(u) \Omega^{(2)}(u) = \langle \mathbf{1}_{12} | , \quad \Omega^{(i)}(u) := \mathbf{R}_{a1}^{(i)}(u) \cdots \mathbf{R}_{al}^{(i)}(u) .$$

It is of particular importance to note the following relation

['04, Beisert, Staudacher].

$$\mathbf{H}_{12} = \frac{d}{du} \ln \mathbf{R}_{12}(u) |_{u=0}$$

- The 1-loop dilatation operator, namely, the **spin chain Hamiltonian** is easily obtained.
- The harmonic R-matrix is used to construct building blocks for the scattering amplitude as Yangian invariant ['13, Ferro , Lukowski , Meneghelli , Plefka , Staudacher] , ['13, Chicherin , Kirschner] , ['14, Broedel , Leeuw , Rosso] .

Summary and prospects

Summary

- We have devolved a new formalism in which the Wick contractions are expressed as a singlet projection.
- Due to the defining property of the singlet, the Ward identities automatically follows.
- For three-point functions in the $SU(2)$ sector, the structure constants are given in terms of the pDWPFs.
- The knowledge of the kinematical dependence greatly helps the analysis.
- The monodromy relations at weak coupling are derived using the crossing and inversion relation, both for the fundamental and the harmonic R-matrix.
- The $1/u$ expansion for them generates non-trivial identities for correlators, including the usual Ward identities.

- A general class of the $SL(2)$ sector using $SL(2)_L \times SL(2)_R$ spin chain.
[Kazama, Komatsu, T.N, to appear]
- More on the monodromy relation.
 \Rightarrow *How they constrain the three-point functions?*
- 1-loop corrections
 \Rightarrow *Is it possible to determine the 1-loop corrections using the symmetries?*